



NON-SINGULAR DEVELOPABLE TRIANGULAR BÉZIER PATCHES

Leonardo Fernández-Jambrina

Departamento de Ciencias Básicas Aplicadas a la Ingeniería Naval

ETSI Navales

Universidad Politécnica de Madrid, Spain


leonardo.fernandez@upm.es

<http://dcain.etsin.upm.es/lfj.htm>



Summary

- We show a characterisation of developable surfaces in the form of Bézier triangular patches.
- Constructions used for rectangular patches are not useful, since they produce degenerate triangular patches.
- Explicit constructions of non-degenerate developable triangular patches are provided.
- Interpolation of a developable triangle between a curve $c(u)$, the last ruling and initial velocity of the other bounding curve $d(u)$.



Previous work

- Construction of developable surfaces of low degree [7-8].
- Construction of developable surfaces in dual space [3,9].
- Constructions grounded on the De Casteljau algorithm [1-2,4-6].


1

Introduction

- Cones and cylinders are easily represented as NURBS surfaces.
- On the contrary, more general developable surfaces are difficult to construct due to the vanishing gaussian curvature condition.
- Solutions for Bézier [1-2], spline [4] and rational [5] developable rectangular patches have been found.
- The simplest require linear relations between the vertices of the control net [1].
- For instance, for Bézier rectangular patches, the control polygons of the bounding curves, $\{c_0, \dots, c_n\}$, $\{d_0, \dots, d_n\}$, are related by

$$(1 - \Lambda)c_j + \Lambda c_{j+1} = (1 - M)d_j + M d_{j+1}, \quad j = 0, \dots, n-1,$$

for fixed Λ , M .



2

Triangular ruled surfaces


- A triangular patch of degree n is defined as

$$c(u, v, w) = \sum_{i+j+k=n} \frac{n!}{i!j!k!} u^i v^j w^k c_{ijk}, \quad u + v + w = 1,$$

for a control net $\{c_{ijk} : i + j + k = n\}$ of $(n+2)(n+1)/2$ vertices.

- The boundary of the patch is formed by the curves at $v = 0$, $u = 0$, $w = 0$.
- We wish a triangular ruled patch between the curves at $v = 0$ and $u = 0$, with control polygons $\{c_0, \dots, c_n\}$ and $\{d_0, \dots, d_n\}$.
- By demanding that constant w curves be straight lines we get

$$c_{ijk} = \frac{ic_{i+j} + jd_{i+j}}{i+j}, \quad c_{00n} = c_0 = d_0.$$




3

Developable Bézier surfaces

- A developable surface bounded by two curves $c(u)$, $d(u)$ has the same tangent plane for all points of each ruling,

$$d'(u) \cdot c'(u) \times (d(u) - c(u)) = 0.$$



- This may be solved in terms of blossoms [4],

$$c[u^{<n-1>}, \Lambda(u)] = d[u^{<n-1>}, M(u)],$$

for some functions $\Lambda(u)$, $M(u)$.


- The constant Λ , M case is the one shown in 2.

4

Degenerate triangular patches

- Unfortunately, the constant case produces just degenerate triangular patches,

$$(1 - \Lambda)c_0 + \Lambda c_1 = (1 - M)d_0 + M d_1 \Rightarrow d_1 = \left(1 - \frac{\Lambda}{M}\right)c_0 + \frac{\Lambda}{M}c_1.$$




- This happens also for general Λ , M unless $\Lambda(0) = 0 = M(0)$.
- The problem is that for $c(u, v) = (1 - v)c(u) + v d(u)$, $d'(0)$ is parallel to $c'(0)$ and therefore to $c_u(0, v)$ for all v .

5

Non-degenerate triangular patches


- We start with a curve $c(u)$ with control polygon $\{c_0, \dots, c_n\}$ and the rulings at both ends, the lines r_0 and r_n . We may fix one end of the ruling, d_0 or d_n , but not both, with a developable surface of constant Λ , M with such boundaries [1].
- For triangles, we begin with $c(u)$ and d_n and the unknown line r_0 to get a developable patch $c(u, v) = c(u) + v \mathbf{v}(u)$, $\mathbf{v} = d - c$.
- We shorten the patch along the rulings, $\tilde{c}(u, v) = c(u) + uv \mathbf{v}(u)$.
- And prescribe the velocity $d'(0) = c'(0) + \mathbf{v}(0) = n(c_1 - c_0) + (d_0 - c_0)$ from which we read the auxiliary vertex d_0 .
- Hence, by this procedure it possible to find at least one triangular developable patch with boundary on $c(u)$ and the ruling $d_n c_n$ fixing the value of $d'(0)$.




6

Other triangular developable patches

- Cylinders bounded by a curve $c(u)$ of degree n and generatrices parallel to a constant vector \mathbf{v} are parametrised as $c(u, v) = c(u) + f(u)\mathbf{v}$ where $f(u)$ is a polynomial vanishing at $u = 0$. The other bounding curve is $d(u) = c(u, 1)$.




- Cones bounded by a curve $c(u)$ of degree n and generatrices parallel to $\mathbf{v}(u)$ may be constructed as $c(u, v) = c(u) + u \mathbf{v}(u)$. The degree of the bounding curve $d(u) = c(u, 1)$ is $n + 1$.



7


Conclusions

- A way of representing ruled triangular Bézier patches in terms of the bounding curves is produced.
- It is shown that usual constructions for designing general developable Bézier surfaces produce only degenerate Bézier triangular patches.
- A procedure for constructing general non-degenerate developable Bézier triangular surfaces is shown.
- Triangular Bézier cylindrical and conical patches can be also constructed.
- Extension to rational Bézier developable triangular patches is possible.



8

References

1. G. Aumann, "Interpolation with developable Bézier patches", *CAGD* **20**, 601-619 (2003).
2. G. Aumann, "Degree elevation and developable Bézier surfaces", *CAGD* **21**, 667-670 (2004).
3. R.M.C. Bodduluri, B. Ravani, "Design of developable surfaces using duality between plane and point geometries", *CAGD* **25**, 621-632 (1993).
4. L. Fernández-Jambrina, "B-spline control nets for developable surfaces", *CAGD* **24**, 189-196 (2007).
5. L. Fernández-Jambrina, (in preparation) 
6. C.-H. Chu, C.H. Séquin, "Developable Bézier patches: properties and design", *CAGD* **34**, 511-527 (2002).
7. W.H. Frey, D. Bindschadler, "Computer aided design of a class of developable Bézier surfaces", *GM Research Publication R&D* **8057** (1993).
8. J. Lang, J., O. Röschel, "Developable $(1, n)$ - Bézier surfaces", *CAGD* **9**, 291-298 (1992).
9. H. Pottmann, H., G. Farin, "Developable rational Bézier and B-spline surfaces", *CAGD* **12**, 513-531 (1995).

9